



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12Q 1/68, C12N 15/10</p>	<p>A1</p>	<p>(11) International Publication Number: WO 92/13100 (43) International Publication Date: 6 August 1992 (06.08.92)</p>
<p>(21) International Application Number: PCT/GB92/00144 (22) International Filing Date: 24 January 1992 (24.01.92) (30) Priority data: 9101757.4 26 January 1991 (26.01.91) GB (71) Applicant (for all designated States except US): MEDICAL RESEARCH COUNCIL [GB/GB]; 20 Park Crescent, London W1N 4AL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : BROOKES, Anthony, Joseph [GB/GB]; 48 Hyde Avenue, Stotfold, Hitchin, Hertfordshire SG5 4JD (GB). PORTEOUS, David, John [GB/GB]; Cherrytrees, 8 Commo Gardens, Edinburgh EH4 9EH (GB).</p>		<p>(74) Agent: KEITH W. NASH & CO.; Pearl Assurance House, 90-92 Regent Street, Cambridge CB2 1DP (GB). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, RU, SD, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US. Published With international search report.</p>
<p>(54) Title: ANALYSIS OF DNA (57) Abstract A method is disclosed of recovering from two mixtures of DNA those sequences which are present in both mixtures (coincident sequences). DNA in a first mixture is produced in single stranded form to which are annealed capture oligonucleotides with known sequences. The first mixture is combined with single stranded DNA in a second mixture. Sequences in the second mixture anneal to homologous sequences in the first mixture. "Captured" homologous sequences are ligated to the capture oligonucleotides and heteroduplex coincident DNA sequences are then recovered.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

- 1 -

Title: Analysis of DNA

Field of invention

This invention concerns analysis of DNA, in particular a method for recovering from two mixtures of DNA only those sequences present in both mixtures, i.e. coincident or shared DNA sequences in the two mixtures.

Background to the invention

When analysing mixtures of DNA, particularly complex mixtures, it would be useful to be able to identify and recover those sequences present in two mixtures, and in recent years attempts have been made to achieve this end. Success so far has been very limited, although a few techniques have been developed which are applicable in limited, restricted situations. For example, a technique has been developed which enables isolation of human inter-alu fragments coincident between overlapping human-rodent somatic cell hybrids.

The present invention aims to provide a more versatile, generally applicable technique for isolating coincident sequences of DNA from two mixtures.

Summary of the invention

According to the present invention there is provided a

- 2 -

method of recovering from two mixtures of DNA those DNA sequences present in both mixtures, comprising treating DNA in a first mixture in a manner comprising the use of restriction enzyme(s) to produce single stranded DNA fragments with added defined flanking sequences flanking the single stranded DNA fragments; annealing capture oligonucleotides to the added defined flanking sequences of the resulting single stranded DNA; treating DNA in a second mixture using the same restriction enzyme(s) to produce single stranded DNA fragments; combining the products obtained from the first and second mixtures and allowing the single stranded DNA fragments to anneal; joining annealed single stranded DNA from the second mixture present as a heteroduplex to the capture oligonucleotides of DNA from the first mixture; and recovering from the resulting mixture sequences captured in the form of heteroduplex coincident DNA including the capture oligonucleotides.

Only DNA sequences present in both the first and second mixtures will form heteroduplex coincident DNA including the capture oligonucleotides, with a sequence from the first mixture carrying the capture oligonucleotides annealing with the coincident sequence from the second mixture. Other DNA fragments will either not anneal or will anneal with other single stranded DNA not including the capture oligonucleotides. The capture oligonucleotides can then be used to recover the heteroduplex DNA of interest, for example by using the polymerase chain reaction (PCR) against the capture oligonucleotides.

The method of the invention, like known methodologies, utilises the formation of heteroduplex species to

- 3 -

distinguish between overlapping and non-overlapping components in two mixtures. However, the known methods utilise two double stranded mixtures of DNA in the formation of this heteroduplex and then achieve its isolation by some means dependent upon pretreatment of the ends of the DNA molecules. In contrast, in the method of the present invention the DNA from the first mixture is initially converted into a single stranded sequence with defined double stranded flanking sequences (by employing capture oligonucleotides). This structure can then act as a sequence specific trap for any related fragments in the unmodified second mixture. Once trapped a joining step, e.g. ligation, is then employed to join the ends of the trapped molecule to the capture oligonucleotides, i.e. specifically to tag the coincident sequences from the second source. By having the modified first source DNA in excess it is possible to drive heteroduplex formation to completion much more readily than is possible in alternative schemes.

Single stranded DNA fragments obtained from the first mixture by treatment with restriction enzymes are conveniently cloned into M13 (or other suitable cloning vehicles) to produce single stranded DNA copies with added defined flanking sequences (from M13 or the other suitable cloning vehicle) to which the capture oligonucleotides are annealed. Other techniques including PCR could also be used for this purpose. The use of PCR has the advantage that it allows for greater flexibility in the choice of the flanking sequences by altering the sequence of the PCR primers.

The capture oligonucleotides may be of any suitable length and are typically about 30-40 base pairs long, with for

- 4 -

example about half of the sequence being designed to bind M13 and the other half being random.

The mixtures are conveniently treated with two restriction enzymes, e.g. EcoRI and PstI, but other combinations are clearly possible.

Depending on techniques used, it may be desirable to functionally isolate any unhybridised capture oligonucleotides before combining the products obtained from the first and second mixtures, and this is conveniently achieved by adding a further oligonucleotide with sequences complementary to the capture oligonucleotides. In particular, such treatment is desirable when PCR is used to recover the coincident DNA of interest.

As mentioned above, the sequences captured in the form of heteroduplex coincident DNA including the capture oligonucleotides may be recovered by PCR. Before using PCR it is necessary to separate these target sequences from the remainder of the first mixture DNA. This may be achieved by a size purification step, e.g. gel electrophoresis. Other techniques may alternatively be used, such as specific modification or degradation of the first mixture sequences achieved by virtue of their single stranded nature.

It may be possible to control the specificity of the method of regulating the degree of homology or identity required for recovery of coincident sequences. The method selects for DNA fragments which are perfectly matched at both ends but a degree of internal mismatch can be tolerated. However, it may be possible to control the

- 5 -

stringency of the method, and by employing techniques such as single stranded modification and/or degradation of annealed heteroduplexes it may be possible for sequences which share up to 100% sequence identity to be recovered by the method of the invention.

The method of the invention is very versatile and generally applicable, and is useful in the analysis of complex mixtures of DNA. For example, the method may be used for isolating coincident sequences between complex sources such as total mammalian genome DNA, type A e.g. Human, and inter-mammalian somatic cell hybrid DNA, sub-AB e.g. a human chromosome or fragment thereof in a complete rodent genome, in the absence of a background of sequences derived from the rodent partner, or from human chromosome regions not represented in the hybrid cell. The method can also be used for isolating highly conserved DNA sequences between distinct species, and for isolating invariant DNA sequences between unrelated individuals from the same species, e.g. regions of linkage disequilibrium spanning disease genes in human populations. The method may also be used for integrating positional information with expression profile, i.e. for isolating candidate genes and exons based on combining available information on map location and tissue restricted expression. The method may also be applicable to microdissection libraries and cDNA libraries. It will be apparent that many other applications are also possible.

The method of the invention is thus of general interest and relevance to all genome mapping, evolution, genetic disease and expression studies.

The invention will be further described, by way of

- 6 -

illustration, in the following Example and by reference to the accompanying figures, in which:

Figure 1 is a schematic representation of one embodiment of the method of the invention applied to two DNA mixtures A and B; and

Figure 2 is a Southern blot analysis of probes 3a and 4a.

Example

DNA sequences common to two mixtures of DNA, mixture A and mixture B, were recovered using the coincident sequence cloning (CSC) method illustrated schematically in Figure 1. Briefly, DNA fragments from mixture A were converted into short, orientated and single stranded molecules with added defined flanking sequences (from M13) at each end by first digesting with two restriction enzymes and then cloning into M13. A pair of synthetic capture oligonucleotides ('capture oligos') was then annealed to this modified form of mixture A (library A). Mixture B was digested with the same restriction enzyme pair that was previously used to process mixture A, and then alkali denatured. Library A and mixture B were combined and allowed to anneal in a reaction driven to completion by the components of library A. Following a ligation step, mixture B sequences were purified from those of library A by preparative alkali agarose gel electrophoresis. Coincident species were then selectively recovered from this material by employing the polymerase chain reaction with primers derived from the capture oligonucleotide sequences. PCR products were finally cloned and analysed.

- 7 -

In this example, DNA mixtures of high complexity were used, with mixture A comprising 1000 fragments of human DNA, and mixture B comprising the total DNA of a somatic cell hybrid called 1W1, which is a human-mouse hybrid with chromosomes 11 and Xpter as the sole human component (see reference 1).

To produce DNA mixture A total human DNA was digested to completion with the enzyme pair EcoRI/PstI and fragments of size 0.1-0.5kb were isolated by preparative agarose gel electrophoresis and cloned into EcoRI/PstI digested M13mp18. 1,000 plaques were picked at random and grown in 150uL cultures. These were pooled for the preparation of single stranded DNA. 5' and 3' capture oligonucleotides (see below) were added to 1ug of this DNA at a molar ratio of 1:1 in 40uL 10mM Tris/HCl, 1mM MgCl₂ pH7.5 and the mixture heated to 65°C and allowed to cool to 37°C over approximately 30 minutes. To bind any unhybridised capture oligonucleotides, oligonucleotide 485 (see below) was then added, in 1uL H₂O, at a molar ratio to each capture oligo of 10:1. This mixture was left at 37°C for 15 minutes and then placed on ice.

15ug of the DNA mixture B was digested with the enzyme pair EcoRI/PstI and then phenol, chloroform and ether extracted and ethanol precipitated. Following resuspension in 50uL H₂O, the sample was denatured by adding 50uL 0.34M NaOH and placing at 37°C for 30 minutes. 100uL of a prechilled 1:1 mixture of 0.34M HCl and 0.1M Tris/HCl pH7.5 was added to neutralise the solution and the sample placed on ice. The modified DNA from the first mixture was then added and the total sample ethanol precipitated. After resuspension in 18uL H₂O both 3uL of 4M NaCl, 50mM EDTA, 0.1M Tris/HCl pH7.8 and 9uL formamide

- 8 -

were added and the mixture allowed to anneal by submerging overnight in a 45°C waterbath. The sample was precipitated and resuspended in 10uL 0.5M EDTA, 5mM Tris/HCl pH7.5. A ligation was then performed at 16°C for 3 hours in 20uL 50mM Tris/HCl, 10mM MgCl₂, 10mM dithiothreitol, 1mM spermidine and 1mM ATP pH7.4 using 0.5 units of T4 DNA ligase.

The ligated DNA was passed through a 1.3% preparative alkaline agarose electrophoretic gel (30mM NaOH, 2mM EDTA buffer) from which single stranded DNA fragments in the 0.1-0.5kb size range were recovered as set of five fractions called F1 - F5. The fractions were recovered as gel slices to be diluted twofold in H₂O and melted at 65°C.

The five fractions were each amplified by PCR using primers derived from the capture oligo sequences in order to recover coincident DNA. 30 cycles of PCR were performed upon 1uL aliquots of the purified DNA using oligos 596/789 (see below). 1uL of these reactions was then further amplified by 22 cycles of PCR using oligos 790/996 (see below).

All PCR reactions were carried out in 50uL PCR buffer (10mM Tris/HCl, 50 mM KCl, 1.5mM MgCl₂, 0.2mM each dATP/dTTP/dCTP/dGTP, 0.01%w/v gelatin, 0.05% each Tween20 and NP40 detergents, pH8.3 at 25°C) using 2 units Amplitaq enzyme and a Hybaid Intelligent Heating Block on mode 2 (plate) control. Denaturing steps were at 99°C for 45 seconds with an extended time of 2 minutes for the first cycle. Extension reactions were done at 74°C for durations of 2 minutes for the first ten cycles, 2.5 minutes for the second ten cycles and 3 minutes for any

- 9 -

further cycles. Annealing steps were of 2 minute duration at 58°C for oligos 596/789 and 52°C for oligos 790/996.

The PCR products were separated on a 1.3% neutral agarose gel. Major bands were excised and recovered by agarose digestion, phenol/chloroform extraction and ethanol precipitation. These products were then cloned following EcoRI/PstI digestion into the pBluescribe plasmid vector, and plated to give product libraries F1 - F5.

To analyse the products of this experiment between 2 and 6 clones were picked at random from each product library. These were examined by cross-hybridisation. In the case of product library F3 two distinct isolates were thus obtained. These were used as probes upon library F3 enabling 14 non-hybridising colonies to be located. From there a further 3 distinct products were subsequently identified. These results are summarised in Table 1.

All distinct products were used to probe Southern blots of EcoRI digested human, mouse and 1W1 DNA in the absence of competitor DNA. In all cases the probes had clearly been derived from human sequences present within 1W1. In the majority of cases a single hybridising band was detected with only 1 case showing hybridisation to a high copy number repeat element. By sequencing each product it was observed that a number of isolates shared a similar sequence even though all had given unique single bands upon Southern analysis. One such product was therefore used as a probe on genomic blots under various stringencies. This family of products were thus shown to have been derived from a low copy repeat element. Sequence database searches were also performed for each coincident sequence cloning product. A summary of these

- 10 -

results is presented in Table 2, in which "Line 1" is human KpnI repeat (reference 2). "L1HEG" is a sequence within a repeat rich region at the beta-globin cluster (reference 3), and "P450c17/SigmaG3" is a sequence upstream of both P450c17 (reference 4) and immunoglobulin heavy C gamma 3 (reference 5) genes. Examples of Southern blot results are shown in Figure 2. Washing stringencies for each experiment are given in the Figure.

All of the products obtained were found to be genuinely coincident indicating a specificity of 100%. Furthermore given the proportion of the human genome present within 1W1 (5%) and the number of mixture A molecules examined (1,000) then the eight products isolated indicate a minimum recovery efficiency of 16%. The spectrum of the coincident species obtained probably reflects their abundance and/or ease of amplification by PCR.

Oligonucleotide sequences

5' capture oligo:

GGACGGGTCGACACGCGAGGAGCCAAGCTTGCATGCCTGCA

3' capture oligo:

AATTCGTAATCATGGTCATAGAGCACCCGTGCTACCGGAACG

485:

TGATTACGAATTGGTGCAGGCATGCAAG

PCR oligos:

596 GGACGGGTCGACACGCGAGG

789 CGTTCCGGTAGCACGGG

790 GCCAAGCTTGCATGCCTG

996 GCTCTATGACCATGATTACG

PRODUCT LIBRARY	NUMBER OF CLONES EXAMINED	DISTINCT PRODUCTS
1	4	1.2a
2	6	
3	3	2a
	14	3a
		3-1, 3-15, 3-20
4	4	4a
5	2	5a

Table 1

CSC products examined and their
product library of origin.

CSC PRODUCT	HUMAN +ve	1W1 +ve	GENOMIC SOUTHERN (high stringency)	DATABASE SEARCH
1.2a	+	+	Repeat	Line 1
2a	+	+	Unique	} LIHEG
3a	+	+	Unique	
3-1	+	+	Unique	
3-15	+	+	Unique	
3-20	+	+	Unique	
4a	+	+	Unique	UNKNOWN
5a	+	+	Unique	P450c17/SigmaG3

Table 2 Summary of result from Southern blot and database search analysis of CSC products. 'Line 1': human KpnI repeat (ref 2), 'LIHEG': sequence within a repeat rich region at the β -globin cluster (ref 3), 'P450c17/SigmaG3': sequence upstream of both the P450c17 (ref 4) and immunoglobulin heavy C γ 3 (ref 5) genes.

- 13 -

REFERENCES

1. Porteous, D.J., Morten, J.E.N., Cranston, G., Fletcher, J.M., Mitchel, A., vanHeyningen, V., Fantes, J.A., Boyd, P.A. and Hastie, N.D. (1986) Mol. Cell. Biol. 6, 2223-2232.
2. Skowronski, J., Fanning, T.G., and Singer, M. (1988) Mol. Cell. Biol. 8(4), 1385-1397.
3. Rogan, P.K., Pan, J. and Weissman, S.M. (1987) Mol. Biol. Evol. 4(4), 327-342.
4. Picado-Leonard, J. and Miller, W.L. (1987) DNA. 6(5), 439-448.
5. Hisajima, H., Nishida, Y., Nakai, S., Takahashi, N., Ueda, S. and Honjo, T. (1983) Proc. Natl. Acad. Sci. USA. 80, 2995-2999.

- 14 -

Claims

1. A method of recovering from two mixtures of DNA those sequences present in both mixtures, comprising: treating DNA in a first mixture in a manner comprising the use of restriction enzyme(s) to produce single stranded DNA fragments with added defined flanking sequences flanking the single stranded DNA fragments; annealing capture oligonucleotides to the added defined flanking sequences of the resulting single stranded DNA; treating DNA in a second mixture using the same restriction enzyme(s) to produce single stranded fragments; combining the products obtained from the first and second mixtures and allowing the single stranded fragments to anneal; joining annealed single stranded DNA from the second mixture present as a heteroduplex to the capture oligonucleotides of DNA from the first mixture; and recovering from the resulting mixture sequences captured in the form of heteroduplex coincident DNA including the capture oligonucleotides.
2. A method according to claim 1, wherein DNA from the first mixture is produced in single stranded form by cloning in M13.
3. A method according to claim 1, wherein DNA from the first mixture is produced in single stranded form by PCR.
4. A method according to any one of the preceding claims, wherein the capture oligonucleotides are in the range of 30-40 bases long.
5. A method according to any one of the preceding claims, wherein an oligonucleotide complementary to the capture

- 15 -


oligonucleotides is added to the products of the first DNA mixture before combining the products of the first and second DNA mixtures.

6. A method according to any one of the preceding claims, wherein heteroduplex coincident DNA is recovered from a mixture by means of PCR.

7. A method according to any one of the preceding claims, wherein the products of the second DNA mixture possess 100% sequence homology with the products of the first DNA mixture to which they anneal.

8. A method according to any one of the preceding claims, wherein the coincident sequence identified by the method is a DNA sequence which is highly conserved between distinct species.

9. A method according to any one of claims 1 to 7, wherein the coincident sequence identified by the method is a DNA sequence which is invariant between unrelated individuals of the same species.





2/2

PROBE 4a

HUMAN	MOUSE	1W1
		

PROBE 3a

HUMAN	MOUSE	1W1
		

HUMAN	MOUSE	1W1
		

kb

← 23

← 9.4

← 6.6

← 4.4

← 2.3

← 2.0

0.1 x SSC,
0.1% SDS,
65°C.


6 x SSC,
0.1% SDS,
65°C.

0.1 x SSC,
0.1% SDS,
65°C.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 92/00144

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 C12Q1/68; C12N15/10		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12Q ; C12N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	WO,A,8 901 526 (GENELABS INC) 23 February 1989 see the whole document ---	1
A	WO,A,8 807 585 (GENELABS INC) 6 October 1988 ---	
A	WO,A,8 912 695 (GENELABS INC) 28 December 1989 ---	
A	EP,A,0 372 524 (THE GENERAL HOSPITAL CORP) 13 June 1990 ---	
P,A	WO,A,9 103 573 (BOEHRINGER MANNHEIM) 21 March 1991 see the whole document ---	1
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
08 APRIL 1992		0 7. 05. 92
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		MOLINA GALAN E. 

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. GB 9200144
SA 55733**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 08/04/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-8901526	23-02-89	AU-A- 2318288	09-03-89
WO-A-8807585	06-10-88	AU-A- 1626288	02-11-88
WO-A-8912695	28-12-89	AU-B- 620628	20-02-92
		AU-A- 3852389	12-01-90
		EP-A- 0419571	03-04-91
EP-A-0372524	13-06-90	AU-A- 4749390	10-07-90
		WO-A- 9006670	28-06-90
WO-A-9103573	21-03-91	DE-A- 3929030	07-03-91
		AU-A- 6271990	08-04-91

THIS PAGE BLANK (USPTO)